## (19) World Intellectual Property Organization International Bureau





### (43) International Publication Date 16 October 2003 (16.10.2003)

## **PCT**

# (10) International Publication Number WO 03/085132 A2

(51) International Patent Classification7:

<del>-----</del>

C12Q 1/68

(21) International Application Number:

PCT/IB03/01791

(22) International Filing Date:

9 April 2003 (09.04.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/370,690

9 April 2002 (09.04.2002) US

(71) Applicant (for all designated States except US): EPIGE-NOMICS AG [DE/DE]; Kastanienallee 24, 10435 Berlin (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): BERLIN, Kurt [DE/DE]; Marienkäferweg 4, 14532 Stahnsdorf (DE).

(74) Agent: SCHUBERT, Klemens; Neue Promenade 5, 10178 Berlin (DE). (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**A2** 

## (54) Title: METHOD FOR ANALYSIS OF METHYLATED NUCLEIC ACIDS

(57) Abstract: A method is disclosed providing analysis of the degree of methylation within nucleic acid samples, including the degree of methylation within CpG islands. After bisulfite treatment of a nucleic acid sample to convert cytosines to uracils, multiple species of paired oliogonucleotide primers and optionally a methylation insensitive reference primer pair are used to amplify target sequences within the sample by methylation specific PCR. Amplification of multiple primer pairs is combined with the use of a real time PCR. Amplificates of primer pairs are detected and quantified by comparison, thus allowing for a detailed, more specific, and quantifiable analysis of the methylation status within a complex CpG methylation pattern of a nucleic acid sample. Primer pairs and a kit are also disclosed.